LSMITHSONIAN MISCELLANEOUS COLLECTIONS,

VOLUME 153, NUMBER 5
(Publication 4753)

Roebling Fund

A LONG-RANGE FORECAST OF TEMPERATURE FOR 19 UNITED STATES CITIES

By

C. G. ABBOT, D.Sc.
RESEARCH ASSISTANT, SMITHSONIAN INSTITUTION

AND

MRS. LENA HILL RESEARCH ASSOCIATE, SMITHSONIAN INSTITUTION



CITY OF WASHINGTON
PUBLISHED BY THE SMITHSONIAN INSTITUTION PRESS
MAY 31, 1969

OF THE
AMERICAN MUSEUM
OF NATURAL HISTORY



SMITHSONIAN MISCELLANEOUS COLLECTIONS VOLUME 153, NUMBER 5

(Publication 4753)

Roebling Fund

A LONG-RANGE FORECAST OF TEMPERATURE FOR 19 UNITED STATES CITIES

By

C. G. ABBOT, D.Sc. RESEARCH ASSISTANT, SMITHSONIAN INSTITUTION

AND

MRS. LENA HILL RESEARCH ASSOCIATE, SMITHSONIAN INSTITUTION



CITY OF WASHINGTON
PUBLISHED BY THE SMITHSONIAN INSTITUTION PRESS
MAY 31, 1969

Roebling Fund

A LONG-RANGE FORECAST OF TEMPERATURE FOR 19 UNITED STATES CITIES

By C. G. ABBOT, D.Sc.
Research Associate, Smithsonian Institution

and

MRS. LENA HILL Research Assistant

INTRODUCTION

The Research findings set forth in this study are the concluding part of an investigation outlined and ordered by Dr. Samuel Pierpont Langley in 1905. Its objects, and the successful progress made in the ensuing 63 years, are briefly but lucidly summarized in Solar Variation, a Weather Element, a paper prepared by C. G. Abbot at the invitation of President Seitz of the National Academy of Sciences, and published in its Proceedings for December 1966.¹ Besides filling volumes 2 to 7 of the Annals of the Astrophysical Observatory of the Smithsonian Institution, explanations and details are contained in the more than 150 papers published by members of the staff of the Astrophysical Observatory during the years 1900 to 1968 in the Smithsonian Miscellaneous Collections. A few are published in other scientific books and periodicals of the United States and foreign countries. A list of the most important of these 150 papers is appended.

It is now three generations since Dr. Langley gave his directive

¹ Reprints of this article are available from the Publications Distribution Section, Smithsonian Institution Press, Washington, D.C. 20560. See footnote 6 for complete citation.

of 1905. A summary of the most important results which flowed from it may be surprising to some of this generation.

- 1. Though Langley deprecated our seeking to determine the exact value of the solar constant of radiation, we had to have the apparatus necessary to obtain it in order to fulfill his other objects. Indeed, as early as 1930, we did obtain a very good value of the solar constant. By 1952, as the average of about 9000 daily values observed from several high mountains, we obtained, from results of the years 1923 to 1952, our published value: 1.946 calories per cm² per minute. During the past year (1967-1968) space observers, observing the sun from outside the atmosphere, have twice obtained 1.95 calories, as published by Dr. A. J. Drummond and associates.² So there is essentially perfect agreement.
- 2. From 1902 to 1914 the staff of the Astrophysical Observatory (then including Andrew Kramer, instrumentmaker, and Messrs. Fowle, Aldrich, and Abbot, observers) designed and constructed about ten instruments for observing solar radiation. These included: the absolute pyrheliometer, four kinds of secondary pyrheliometers, the pyranometer, the two-mirror coelostat, the perfected vacuum-bolometer, and several devices for the spectro-bolometer, and for reducing solar measurements.³
- 3. From 1905 to 1920 about six months each year were spent in solar radiation work on Mt. Wilson, measuring the atmospheric transmission, and computing the solar constant, by Langley's "long method." H. H. Clayton—making high, medium, and low groups of our results—proved by 1916 the sun's radiation to be a controlling world weather element.
- 4. In 1918 a new station was established at Calama, in the nitrate desert of Chile, to observe the sun's radiation daily throughout the year. Clayton found a close correlation between Mt. Wilson and Calama, though situated in opposite hemispheres. But we needed daily measures from a pair of mountain stations, and a solar constant method so quick as to avoid changes of atmospheric transparency. John A. Roebling's generosity enabled us to occupy Mt. Harqua Hala (5672 feet) in Arizona, and also Mt. Montezuma (9000 feet) near Calama throughout several years beginning in 1920.
 - 5. A. F. Moore at Calama, making daily measures of radiation

² Eppley Laboratory, Reprint Series No. 33, 1967.

³ See C. G. Abbot, Solar Variation and Weather—A Summary of the Evidence, Completely Illustrated and Documented, Smithsonian Miscellaneous Collections, volume 146, No. 3 (Publication 4545).

from a zone of sky near the sun with the pyranometer, laid the foundation for the "short method" of determining the atmospheric spectral transparency in 40 wavelengths. The "short method" for the solar constant was perfected in 1923, and applied to measure the solar constant daily for 9000 days at several high mountain stations over a period of 30 years, from 1923 until 1952.4

- 6. Meanwhile, Dr. George E. Hale, at Mt. Wilson, discovered the cycle of 22 years 9 months in the magnetic fields of sun spots. Our 30-year record of daily solar constant measures plainly revealed a corresponding master cycle of 273 months in solar variation. It has an amplitude of 3 percent and, like music, has many exact subordinate harmonics. We discovered 27 such harmonics in solar variation, all exact fractions of 273 months. We found also non-periodic trends, up and down, for the sun's radiation. These opposing trends, occurring about twice a month, and with amplitudes of approximately 1 percent, were found to be an important cause of temperature changes.
- 7. Following Dr. Langley's prevision, we found all of the 27 regular solar harmonics, and also all of the nonperiodic trends, to be plainly effective in weather.
- 8. Finally, Langley's hope for long-range weather predictions also are confirmed. In five publications in the Smithsonian Miscellaneous Collections series ⁵ forecasts of precipitation at 55 cities on six continents were tabulated for as much as two generations in advance. In publication 4711, identified below in footnote 5, and in this publication, long-range forecasts are also tabulated for 30 cities in the United States. In Solar Variation, a Weather Element, ⁶ Figure
- ⁴ See C. G. Abbot, Forecasting from Harmonic Periods in Precepitation, Smithsonian Miscellaneous Collections, volume 148, No. 8, 1966 (Publication 4659).
- ⁵ See C. G. Abbot, *Sixty-Year Weather Forecast*, Smithsonian Miscellaneous Collections, volume 128, No. 3, 1955 (publication 4211).
- C. G. Abbot, A Long-Range Forecast of United States Precipitation, Smithsonian Miscellaneous Collections, volume 139, No. 9, 1960 (Publication 4390).
- C. G. Abbot, *Precipitation in Five Continents*, Smithsonian Miscellaneous Collections, volume 151, No. 5, 1967 (Publication 4694).
- C. G. Abbot, Supplement to a Long-Range Forecast of United States Precipitation (Smithsonian Publication 4390), Smithsonian Miscellaneous Collections, volume 152, No. 5 (Publication 4711).
- C. G. Abbot, Solar Magnetism and World Weather, Smithsonian Miscellaneous Collections, volume 152, No. 6, 1967 (Publication 4722).
- ⁶ C. G. Abbot, Solar Variation, a Weather Element, Proceedings of the National Academy of Sciences, volume 56, No. 6, pages 1627-1634, December 1966.

2 shows large average effects on the temperature of Washington from rising and falling trends of solar radiation in all months of the year.

TEMPERATURE FORECASTS FOR 19 UNITED STATES CITIES

The temperatures forecasted here for 19 United States cities were computed by us from electronic tabulations by Jonathan Wexler, and were prepared by him from World Weather Records, 1880 through 1949. Excepting Nashville, Tennessee, all our long-range predictions extend from 1967 to 1972. For Nashville we predicted from June 1942, so that we could show the effect of bombing-both with uranium preparations, and later with hydrogen bombs-from 1944 to 1964.

TABLE 1.—Cities Where Temperature is Forecasted

Bismarck, North Dakota
Charleston, South Carolina
Chicago, Illinois
Cincinnati, Ohio
Denver, Colorado
Eastport, Maine
El Paso, Texas

Galveston, Texas Helena, Montana Little Rock, Arkansas Marquette, Illinois Mobile, Alabama Nashville, Tennessee Washington, D.C.

Phoenix, Arizona Portland, Oregon Sacramento, California Sante Fe, New Mexico Spokane, Washington

Table 2.—Normal Monthly Temperatures, in Fahrenheit, for 19 United States Cities, Means, 1880-1949

A. Sunspots >20. B. Sunspots <20.

	Віѕм	ARCK	CHARL	ESTON	Сни	CAGO	CINCINNATI		
	A	В	A	В	A	В	A	В	
January	10.5	8.5	50.9	49.8	27.2	25.0	33.8	32.3	
February	10.4	11.2	51.9	51.2	27.6	27.6	34.2	35.0	
March	23.3	26.4	56.5	57.7	35.6	37.5	42.4	44.8	
April	44.7	42.8	63.9	64.6	48.0	47.2	54.6	54.4	
May	55.9	54.5	72.7	72.1	59.2	57.8	65.6	63.9	
June	65.6	64.0	78.5	78.1	68.8	67.6	74.0	72.8	
July	70.8	77.1	80.7	80.1	74.6	73.5	78.3	76.9	
August	67.9	68.7	79.9	80.0	73.0	73.0	75.4	75.1	
September	57.5	58.7	76.7	76.4	65.7	66.7	68.7	69.4	
October	46.2	44.3	67.3	67.5	55.4	54.8	57.8	57.7	
November	29.6	27.7	58.0	58.2	42.3	40.6	45.6	44.7	
December	16.4	15.6	52.0	50.8	31.1	29.5	36.6	35.3	

	DEN	IVER	EAST	PORT	EL I	PASO	GALVI	ESTON
	A	В	A	В	A	В	A	В
January	32.1	29.9	21.2	21.4	44.5	43.8	54.9	53.3
February	31.5	32.9	21.6	22.1	49.0	49.4	56.9	55.8
March	37.6	39.4	29.8	29.6	55.2	55.7	60.9	62.2
April	48.5	47.8	38.8	37.6	63.4	63.5	68.6	68.5
May	57.5	56.5	47.4	47.0	72.6	71.8	75.1	74.8
June	68.3	66.5	54.6	54.4	80.5	80.5	80.5	81.0
July	73.2	72.7	59.9	59.5	81.3	81.6	82.7	82.9
August	71.7	71.3	59.7	60.1	80.6	79.3	83.1	80.0
September	62.5	63.2	55.6	55.6	74.5	74.3	79.9	73.1
October	51.6	51.0	47.8	47.6	64.6	64.0	72.7	63.1
November	40.9	38.9	37.1	37.4	52.5	51.7	64.2	56.5
December	32.3	32.8	25.5	26.6	44.7	45.0	56.7	
	HEL	ENA	LITTLE	Rock	Marg	UETTE	Mon	BILE
	A	В	A	В	A	В	A	В
January	21.9	19.5	43.0	40.6	17.8	16.8	52.2	51.0
February	23.2	23.9	44.2	44.7	16.1	17.3	54.7	53.7
March	31.2	32.8	51.2	53.7	23.8	26.5	58.6	60.2
April	41.6	43.5	62.3	62.2	38.4	38.4	66.4	66.5
May	52.7	52.2	70.1	69.4	50.0	48.1	73.5	73.2
June	60.4	59.6	77.9	77.3	59.5	58.5	79.3	79.4
July	68.2	68.5	81.0	80.5	65.3	65.1	80.5	80.3
August	66.2	67.2	80.0	79.6	63.7	64.0	80.3	80.3
September	55.4	55.8	74.0	74.4	56.9	57.6	77.6	77.5
October	45.8	44.6	63.3	63.5	47.4	46.2	68.0	68.8
November	33.7	32.4	52.8	51.6	34.1	32.9	58.9	58.5
December	23.8	25.3	43.6	43.9	23.1	23.0	53.4	52.3
	SACRA	MENTO	Nash	VILLE	Рно	ENIX	Port	LAND
	A	В	A	В	A	В	A	В
January	45.4	45.7	40.2	38.0	51.0	50.8	39.7	38.5
February	50.1	50.5	41.1	41.1	54.8	55.1	41.8	42.2
March	55.0	54.2	47.7	50.8	60.7	60.4	46.8	46.9
April	58.3	59.0	58.8	59.2	67.3	67.8	51.1	51.9
May	63.9	63.9	68.6	67.5	76.0	75.6	57.5	57.5
June	69.2	70.5	76.4	75.4	84.8	84.6	61.8	62.0
July	73.8	74.3	79.4	78.3	90.5	90.0	66.9	67.2
August	72.9	73.5	77.6	77.7	88.7	88.3	66.6	67.3
September	70.8	70.0	72.0	72.4	83.4	82.5	61.7	61.8
October	63.3	63.0	60.6	61.2	71.4	70.6	54.7	53.9
November	54.1	53.7	49.7	48.8	60.1	58.8	46.4	46.1
December	46.2	46.5	41.3	41.5	52.1	52.3	41.0	41.7

	SANT	E FE	Spok	ANE	Washington, D. C.				
	A	В	A	В	A	В			
January	29.1	28.7	30.1	26.0	35.9	33.9			
February	32.8	33.0	30.4	31.6	35.3	35.6			
March	39.0	39.2	39.5	39.8	42.5	44.5			
April	47.6	47.5	48.4	48.0	53.8	54.5			
May	56.5	55.6	56.1	56.0	65.0	64.0			
June	66.8	65.3	62.4	62.4	72.9	72.4			
July	68.9	68.6	69.7	70.2	76.9	76.3			
August	67.7	67.2	68.6	68.9	74.6	74.8			
September	61.2	61.5	58.8	59.1	68.7	68.6			
October	50.6	50.2	48.9	48.0	57.3	57.5			
November	39.2	38.3	37.6	36.9	46.6	46.7			
December	30.2	30.4	30.6	30.5	37.1	37.0			

As described in several publications listed in the appendix, all of our long-range forecasts, both precipitation and temperature, are compiled by adding the values obtained to represent the weather effects of 27 regular periods, all exact harmonies of 273 months. Hence the forecasts are not *simple* monthly values, but *complex smoothed monthly values*. To compare with them *fairly*, the observed monthly values must also be *smoothed*. The following Tables 3 and 4 give the forecasts, the directly observed monthly values, and the monthly observed values *smoothed* by three-month consecutive smoothing. The differences tabulated in Table 3 are between the monthly forecasts and the three-month smoothed observed temperatures.

THE EFFECTS OF ATOMIC AND HYDROGEN BOMBS ON FORECASTS AT NASHVILLE

Figure 6 in Solar Variation, a Weather Element * shows graphically how prejudicially the atomic bombing in Japan, and the later hydrogen bombs exploded by the United States and Russia in the Pacific and Arctic Oceans, affected long-range forecasts of precipitation at Tokyo and Lagos. Many similar scatter-graphs of long-range precipitation forecasts at stations in distant regions are on file at the Smithsonian. We wish now to show, in another kind of graph, what effect atmospheric bombing explosions appear to have produced on long-range temperature forecasts in central United States.

⁷ Ibid.

Figure 1 graphs the march of long-range temperature forecasts at Nashville, Tennessee, from 1942, before atomic bombs were made, through 1965, after the United States and Russia had exploded mighty hydrogen bombs at intervals from 1949 to 1960.




FIGURE 1.—Nashville temperature departures.

nning Means	HELENA		-1.3	-4.7	-2.1	-3.5	-3.2	3.3	-0.5	-3.2	-2.1	+0.3	+0.6	+3.3		+2.9	+0.5	+1.1	+2.5	+3.1	+3.0	+1.5	+2.0	+1.0	+1.5	-1.6	-3.0		4.0—	-0.4
TABLE 3.—Forecast of Temperature Departures from Normal 1968 Through 1972 from Three-month Running Means	GALVESTON		-1.5	-2.3	-1.4	-2.2	7.0—	+0.9	+0.1	+0.3	+0.5	+1.0	-0.3	+0.8		+0.9	+0.8	-0.4	+0.4	6.0—	+0.3	+0.9	0.0	+0.9	+0.9	+2.4	-0.3		+1.9	+2.0
972 from Th	Er Paso		+1.5	+1.1	+1.0	-0.1	+1.0	+0.8	+0.2	-0.5	-0.2	0.7	-1.7	-0.3		+1.1	+1.5	+1.5	+2.1	+2.4	+2.1	+1.8	+0.8	+0.4	-0.1	-0.4	-1.0		-2.6	0.0
8 Through 1	EASTPORT		+0.8	+1.6	+0.1	-1:1	-1.4	-1.7	-3.4	-4.1	-4.1	-2.3	-1.7	-1.4		-1.2	8.0—	+0.5	+1.1	-0.1	+0.3	+2.6	+2.2	+2.2	+3.5	+2.8	+4.0		+2.9	+0.5
Normal 196	Denver		+1.7	+0.5	+0.9	-0.5	+1.2	+1.1	+1.4	+2.4	+1.8	+3.6	+1.9	+3.8		+1.0	+1.3	-0.2	+0.1	8.0—	-1.6	-2.5	-3.1	-3.3	-3.8	-2.3	-5.2		6.4-	-3.1
artures from	CINCINNATI		+0.9	+1.0	+3.5	+3.4	+4.0	+4.8	+3.5	+3.0	+1.5	+2.3	+1.0	+1.9		+1.0	+2.3	+1.2	+0.8	+1.3	+2.3	+2.1	+3.6	+4.8	+3.4	+3.1	+2.8		+2.2	+2.2
erature Dep	CHICAGO		+1.4	6.0—	+1.0	+0.4	+0.9	-0.7	-0.1	9.0—	-1.8	+0.3	-1.3	+1.0		+1.9	+2.1	+1.7	+3.3	+3.9	+4.2	+3.4	+4.0	+5.6	+4.3	+3.4	+2.7		+2.9	+3.5
ecast of Temp	CHARLESTON		-1.8	-2.7	-2.4	-2.2	-1.6	6.0—	-0.2	+0.4	-1.4	-1.4	-1.5	-1.8		-1.3	-1.4	-1.6	-1.5	+1.5	9.0—	+0.7	+2.0	+3.0	+3.3	+2.9	+4.5		+4.8	+4.3
ABLE 3.—For	BISMARCK		-2.5	-3.4	-3.1	-3.3	-2.4	-2.8	-1.2	-2.3	-2.9	+1.3	+1.7	+1.9		+2.5	+1.8	+1.3	-0.2	-1.2	-1.5	6.0—	+0.8	-1.6	7.0—	-1.7	-2.2		8.0—	+1.0
T		1968	January	February	March	April	May	June	July	August	Sepetmber	October	November	December	1969	January	February	March	April	May	June	July	August	September	October	November	December	1970	January	February March

1.0 + 4.5 + 4.5 + 4.5 + 4.5 + 4.5 + 4.5 + 4.5 + 5.1 + 5.1	++++++++++++++++++++++++++++++++++++++	+ + + + + + + + + + + + + + + + + + +
++1.8 ++2.5 ++2.6 ++2.9 ++2.7 ++2.7	++++++++++++++++++++++++++++++++++++++	+0.9 +1.1 -1.0 -1.3 +0.6 +2.3 +2.0
-0.6 -0.7 -0.7 -1.3 -1.3 -1.3 +3.1 +3.2 +3.4 +3.4	++++++++++++++++++++++++++++++++++++++	++++++++++++++++++++++++++++++++++++++
+ + + + + 2.1 + + + 0.2 + 0.2 - 2.0 - 2.0 - 3.0	- 2.7 - 2.8 - 2.1 - 2.1 - 2.1 - 1.5 - 4 - 4 - 4 - 6.1 - 1.5 - 6.1 - 1.5 - 6.1 - 6.1	+ + 2.5 + 2.5 + 2.5 + 2.5 + 1.6 + 1.8 + 0.7
-2.1 -1.2 -1.2 -1.4 -1.3 -1.3 -1.3 -1.3 -1.3 -1.3 -1.3 -1.3	+ + + + + + + + + + + + + + + + + + +	++++++++ 9,6,7,4,4,4,4,4,4,4,4,4,4,4,4,4,4,4,4,4,4
++1.2 0.2 2.4 2.0	1.5.0 1.	- 4.6 - 3.0 - 2.1 - 2.0 - 1.2 - 0.1 - 0.1 + 3.1 + 2.8
+++++5:5 1.1 + + + + + + + + + + + + + + + + + +	+ + + 0.6 + 0.5 + 0.6 + 0.5 + 0.4 + 0.5 - 0.4 -	-3.5 -3.0 -0.5 -1.4 -2.0 -2.4 -0.4 -0.3
++++++++ 4.4.6.5.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2	- 3.3 - 2.7 - 2.7 - 1.9 - 1.1 - 1.6 - 1.1 - 1.0 - 1.1 - 1.1	+ 0.5 - 0.5 - 0.5 - 0.5 - 0.1 + 0.6 + 1.7 + 2.5
+ + + + + + + + + + + + + + + + + + +	- 3.8 - 2.1 - 2.1 - 2.8 - 4.4 - 5.0 - 5.0	+ 2.8 + 6.3 + 6.3 + 8.0 + 8.6 + 10.7 + 110.2
April May June July August September October November December	1971 January February March April May June July August September October November	January February March April May June July August

Table 3.—continued

WASHING- TON, D. C.		-0.1	+0.1	+0.3	+0.7	+1.1	+0.9	9.0—	9.0—	-2.2	-1.7	8.0—	-1.1		-0.3	+0.0	9.0—	-0.1	+1.4	8.0—	+0.5	+0.8	+1.2	+0.9	+0.9	+1.8		+1.5	+1.3	+0.8
SPOKANE		+4.7	+4.1	+5.2	+5.7	+5.9	+6.2	+4.8	+3.1	+0.9	+2.3	+0.4	+0.5		+0.6	6.0—	6.0—	-0.3	0.0	+0.4	+0.6	+0.6	+1.7	+0.4	6.0—	+0.2		-0.3	-1.2	-2.2
SANTA FE		-1.0	-3.2	+0.1	-1.6	-0.7	-1.1	-0.1	-0.3	+0.7	0.0	9.0—	+0.5		+0.6	+0.6	-0.3	+0.6	+1.4	+1.8	+2.2	+2.5	+3.0	+2.0	+1.9	+1.2		+1.2	+2.6	+3.3
PORTLAND		-1.3	+0.9	-0.1	+0.6	+1.1	+0.9	+1.4	+1.2	+1.3	+1.6	+1.0	+1.7		+0.9	+0.6	9.0—	-1.1	-1.5	-1.6	9.0—	-1.0	-0.5	8.0—	+0.2	-0.3		-0.3	-0.1	+0.3
PHOENIX		-0.2	-1.8	-1.2	+0.9	+0.3	-0.3	-1.1	+2.6	+1.3	+2.0	+2.2	+2.0		+1.8	+1.9	+2.5	+1.4	+1.9	+3.0	-0.1	+0.1	6.0—	+0.4	-1.0	-1.2		+0.1	+1.9	+0.3
NASHVILLE		-0.2	-1.2	-2.4	-2.1	-2.4	-3.1	-4.5	-5.0	-5.2	-5.5	6.4-	-3.9		-1.4	-1.6	-2.1	-1.3	-0.3	+0.4	+0.5	+1.8	+2.2	+3.0	+1.0	+2.1		+2.0	+2.5	+0.9
SACRAMENTO		0.0	+0.5	+0.1	+0.3	-0.2	+0.2	7.0—	-2.0	-0.5	+0.9	+0.9	+1.2		+1.9	+3.6	+3.6	+3.8	+3.3	+2.7	+1.8	+1.0	-0.3	+0.3	7.0-	9.0—		-1.0	-1.2	-0.1
Mobile		+0.7	9.0—	+1.0	+0.3	+0.5	+0.7	-0.4	7.0—	-1.9	-1.8	-1.9	-1.7		-1.8	-1.4	-1.6	-1.2	8.0—	+0.5	+1.1	+0.9	+1.1	+1.2	+1.2	+1.8		+1.5	+1.8	-0.1
MARQUETTE		-1.1	+1.0	+1.2	+1.0	+2.0	+2.0	+2.1	+0.6	8.0—	+1.8	0.0	+3.5		+2.8	+2.6	-0.1	+1.5	+0.3	+2.7	+0.3	+2.3	+2.2	+4.5	+1.6	+1.6		+3.1	+3.7	+3.0
LITTLE ROCK		7.0-	-2.4	-0.5	0.0	+0.5	+1.4	+1.3	+1.2	4.0—	+1.3	-0.4	+1.4		+1.1	+0.8	-0.2	+0.9	+1.3	+1.2	+2.1	+2.8	+3.9	+3.3	+3.1	+2.3		+1.4	+2.3	+0.7
	1968	January	February	March	April	May	June	July	August	September	October	November	December	1969	January	February	March	April	May	June	July	August	September	October	November	December	0261	January	February	March

++++++++++++++++++++++++++++++++++++++	1.6 -0.8 -0.8 -1.7 -1.7 -1.4 -2.9 -3.8 -3.8 -2.8 -2.8 -2.8 -3.0	1.3 -1.6 -0.5 -0.9 +1.4 +1.4 +2.9 +2.9 +3.2
3.5 6 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	2.5 - 2.5 - 1.5 - 1.5 - 1.5 - 1.1 - 2.1 - 1.2 - 1.3 - 1.3 - 1.9 - 1.9 - 1.9 - 1.9 - 1.9 - 1.9 - 1.9 - 1.0 -	++++2.5 +++5.1 ++6.7 +6.6 +6.6 +6.6
++++++++++++++++++++++++++++++++++++++	+ + + + 2.1 + + 1.6 + 0.2 - 1.1 - 1.5 - 2.3 - 2.3 - 2.3	-2.4 -1.6 -1.6 -1.0 -1.0 -1.0
-0.7 -0.1 -0.1 -0.2 -0.2 -0.4	1.1 0.0 0.0 0.0 0.0 0.0 0.0 0.0	+ + + + + + + + + + + + + + + + + + +
+1.8 +2.5 +2.3 +1.7 +1.7 +1.0 +2.7 +1.6	+ + + + + + + + + + + + + + + + + + +	-1.0 -1.0 +2.5 +2.5 +2.9 -1.0 -1.0
+ 1.0 + 0.3 + 0.7 + 0.7 + 0.4 + 0.4 - 0.6 - 1.7 - 2.6	- 2.9 - 1.0 - 1.0 - 1.0 - 2.3 - 4.0.1 - 4.0.9 - 4.2.2 - 4.3.3	+ + + + + + + + + + + + + + + + + + +
- 0.8 - 0.2 - 0.5 - 0.3 - 0.3 - 0.5 - 0.5	0.5 0.6 0.8 0.8 0.6 0.6 0.6	+ 0.2 + 1.3 + 2.8 + 2.6 + 2.0 + 1.5 - 0.5
0.4 -0.4 -0.6 -0.9 -1.0 -1.7 -1.7	- 0.9 - 0.9 - 0.9 - 0.9 - 0.9 - 0.6 - 0.6 - 0.6 - 0.9 - 0.0 - 0.0	+ + + + + + + + + + + + + + + + + + +
++++++++++++++++++++++++++++++++++++++	7.2 4 6 4 7 9 7 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	- 4.1.1 - 0.9 - 0.9 - 0.9 - 0.9 - 1.1.1 + 1.1.2
0.5 + 0.1 - 1.3 - 1.6 - 1.6 - 1.7 - 1.7	1.6 1.6 1.6 1.6 1.6 1.1 1.1 1.1 1.0 1.0 1.0 1.0 1.0 1.0 1.0	++++++++++++++++++++++++++++++++++++++
April May June July August September October November December	J971 January February March April May June July August September October November	1972 January February March April May June July Nuggust September

Table 4.—Four-month Mean Temperatures, 1965-1972 All based on three-month running mean data

	Bisn	IARCK			CHAR	LESTON		CHICAGO				
1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	1 965	Pre- dicted	Ob- served	Δ	
I II III		-3.3 -4.2 0.0		I			-3.9 -4.7	I		-0.6 -1.8 -1.4		
1966 I II III	-0.6	+1.6 -3.2 -1.6	-2.6	1966 I II III	+0.9			II	-3.6	0.7 0.9 0.2	+2.7	
		+0.8 -4.4		II		+1.8 -1.8	+0.3 +1.2			+1.0 -1.0		
II	-3.1 -2.2 +1.1				-2.3 -0.6 -1.5			II	+0.5 -0.1 -0.4			
II	+1.4 -0.7 -1.6				-1.4 +0.9 +3.4				+2.2 +3.9 +4.0			
	+0.2 +0.3 -0.2			1970 I II III					+2.5 +1.2 +0.4			
II	-3.5 -3.7 -0.1			II	-2.3 -0.9 +0.8			II	+0.2 -1.4 -4.4			
	+5.8 +10.6			1972 I II III	-0.3 +0.4				-2.1 -1.0			

TABLE 4.—continued

	Cinci	INNATI			DEI	NVER		EASTPORT				
1965 I	Pre- dicted	Observed —1.5	Δ	1965 I	Pre- dicted	Ob- served —1.6	Δ	1965 I	Pre- dicted	Ob- served +2.5	Δ	
III		-3.5 -2.1		III		-3.4 +1.1	-3.0 -1.9		+2.4 +2.5			
1966 I II III	-2.1 -2.5 -1.7	-0.9		II	-0.6		-2.4 -1.0 +4.6			+2.9	-0.1 + 1.8 + 0.1	
1967 I II III	2.8	+0.1 -1.0 -1.8	+1.8	II	+0.8		+4.9 -5.7					
1968 I II III	+2.2 +3.8 +1.7			1968 I II III	+0.6 +1.5 +2.8			1968 I II III	+0.4 2.6 2.4			
	+1.3 +2.3 +3.5			1969 I II III				1969 I II III	1			
1970 I II III	+1.9 -0.6 -2.2			1970 I II III	-3.2 +0.3 +0.7			1970 I II III	+2.0 +1.0 -2.0			
1971 I II III	-3.5 -4.8 -5.0			1971 I II III	-0.2 +0.3 +2.4			1971 I II III	-2.6 -1.7 +0.6			
1972 I II III	-2.9 +0.7			1972 I II III	+5.4 +4.6			1972 I II III	+2.6 +1.6			

TABLE 4.—continued

	El]	Paso			GALV	ESTON		Helena				
1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	
	-1.7 + 1.1	+0.2 -0.6 -1.6		I II III	+2.4		-2.0 -2.2			+1.6 -3.3 $+0.7$		
1966 I II III	+2.3	0.6 0.9 0.4	-2.2	II	+0.8		-3.1 -2.0 +1.5	II			+1.6 -3.0 +4.9	
II		+1.0 -1.3		II			+4.6 -2.2	II				
II	+0.9 +0.4 -0.7			1968 I II III				II	-2.9 -2.6 +0.5			
II	+1.6 +1.8 -0.3			1969 I II III	+0.1				+1.8 +2.4 -0.5			
II	-1.0 +0.6 +3.3			1970 I II III	+2.4			1970 I II III	-0.6			
II	+3.8 +1.9 +1.2			1971 I II III	+3.3 +3.1 +2.1			II	+1.4 +0.3 0.0			
1972 I II III	+3.3 +5.2			1972 I II III	-0.1 +0.6			1972 I II III				

TABLE 4.—continued

	LITTL	в Роск			Marg	QUETTE		Mobile				
1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	
III II	+3.1 +1.5	-0.2 +0.4 +1.0		II II	+0.7 +0.8		2.5 0.7	II III	+1.0 -0.3	+1.4 +0.6 +2.2	-0.4 +2.5	
1966 I II III	-1.6	0.0 -0.8 -0.6	+0.8		+0.3	+0.1	+4.2 -0.2 +1.6		-0.8 -1.2 -0.5	-0.1 +0.6 +1.3	+0.7 +1.8 +1.8	
1967 I II III	+1.0 -1.0 -3.0	+1.9 -2.0			-1.0 -3.0 -0.9		+2.4 +1.8	1967 I II III			,	
1968 I II III	-0.9 +1.1 +0.5			1968 I II III	+0.5 +1.7 +1.1			1968 I II III	+0.4 0.0 -1.8			
1969 I II III	+0.6 +1.4 +3.2			1969 I II III	+1.7 +1.4 +2.5			1969 I II III	-1.5 +0.4 +1.3			
1970 I II III	+1.0 -1.3 -1.6			1970 I II III	+3.1 +2.4 -0.6			1970 I II III	+0.7 -0.6 -1.7			
1971 I II III	-0.8 +1.0 +2.0			1971 I II III	-4.3 -6.2 -5.0			1971 I II III	+0.2 +1.4 +1.4			
1972 I II III	+2.8 +2.6			1972 I II III	-3.6 +0.4			1972 I II III	+1.2 +1.4			

TABLE 4.—continued

	Nashville				Рно	ENIX		PORTLAND			
1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ
I		-0.3		I		-1.9		I		-0.1	
II		-0.7		II			-3.5		-1.0		
III	+2.7	+0.8	-1.9	III	-0.2	+0.7	+0.9	III	-0.9	+1.5	+2.4
1966				1966				1966			
I	+3.6		-3.2	I			+1.8	I		,	+2.1
III	+1.7	-1.8 -0.8	3.5 2.2	III	-1.4 -2.1		+2.6 +1.8	III	-0.2 -0.3		
	T 1.7	-0.0			- L.	-0.0	71.0		-0.0	-[-1,-1	Ţ1.,
1967 I	112	120	1 1 7	1967	1.0	102	120	1967	20	106	126
II		+2.9 -2.7	-3.8			+0.2 -1.5			-2.0 -1.5	+0.6 +1.4	
III	+1.1	2.,	- 0.0	III		1.0	0.1	III		1	,
1968				1968				1968			
	-1.5			I	-0.6				0.0		
II	-3.8			II	+0.4			II	-1.2		
III	—4.9			III	+1.9			III	+1.4		
1969				1969				1969			
I	-1.6			I	+1.9			I	-0.1		
II	+0.6			II	+1.2			II			
III	+1.8			III	— 0.7			III	-0.4		
1970				1970				1970			
I				I	+1.0				-0.2		
II	+0.4			II	+2.2			II	+0.8		
III	-1.6			III	+1.8			III	+0.1		
1971				1971				1971			
	-1.9			I				I	-0.6		
III	-1.2 + 1.8			III	+0.6 -1.9			III	+0.7 +3.5		
	[-1.0				-1.7				F-0.0		
1972 I	+3.8			1972 I	106			1972 I	+3.6		
II	+2.7			II	+0.6 +0.9			II	+1.2		
III	,,			III	1 0.5			III	1		

TABLE 4.—continued

	SACRAMENTO				SAN	ra Fe			Spokane			
1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	1965	Pre- dicted	Ob- served	Δ	
I		0.0		I		+7.6		I		1.5		
II			-0.2		+1.0	•	+3.8		-4.5		+0.5	
III	+1.4	0.0	-1.3	III	+1.6	+6.7	+5.1	III	-2.7	+0.8	+3.5	
1966				1966				1966				
I	+1.8	+0.2	-1.6	I	+1.6		+3.7	I	0.5	+0.6	+1.1	
II	-0.5	,	+1.2		+1.2		+6.0		+1.0		-3.5	
III	-3.5	+1.3	+4.8	III	+0.1	+6.7	+6.6	III	-2.6	+1.6	+4.2	
1967				1967				1967				
I	2.3	-2.4	0.1	I	0.8	+7.0	+7.8	I	-5.2	+1.4	+6.6	
II	+1.0	+0.7	0.3	II		+9.4	+8.6	II		-0.7	+0.4	
III	+1.1			III	0.5			III	+2.1			
1968				1968				1968				
I	+0.2			I	-1.4			I	+4.9			
II	0.6			II				II	+5.8			
III	+0.6			III	+0.2			III	+1.0			
1969				1969				1969				
I	+3.2			I	+0.4			I	-0.4			
II	+2.2			II	+2.0			II	+0.4			
III	-0.3			III	+2.0			III	+0.4			
1970				1970				1970				
I	0.8			I	+2.3			Ι	-1.7			
II	+1.2			II	+3.8			II	-3.3			
III	+0.7			III	+3.2			III	-3.7			
1971				1971				1971				
I	0.8			I	+1.8			I	-2.4			
II	-1.3			II	-0.9			II	+0.7			
III	-1.7			III	-2.2			III	+2.4			
1972				1972				1972				
I	+1.5			I	-1.7			I	+4.4			
II	+2.0			II	0.9			II	+6.5			
III				III				III				

TABLE 4.—continued

	Wash	INGTON	, D. C.
1965	Pre- dicted	Ob- served	Δ
III	+2.5 +0.8	$+1.0 \\ +0.1 \\ +2.9$	-2.4 +2.1
1966 I II III	-0.8 -0.7 -0.3	+3.0 +1.0 +1.9	+3.8 +1.7 +2.2
1967 I II III	-0.8 -1.8 -1.7	+2.3 -0.8	+3.1 +1.0
1968 I II III	+0.2 +0.2 -1.4		
1969 I II III	+0.1 +0.5 +1.2		
1970 I II III	+1.2 +1.2 +1.7		
1971 I II III	-1.3 -3.0 -3.2		
1972 I II III	-1.1 + 1.3		

It is plain that the graph comprises intervals of good forecasts and intervening intervals of distorted forecasts. These intervals may be tabulated as follows:

Period of good forecasts	Number of months	Mean observe minus predict	ed s
June 1942 to November 1944	30	20	
April 1947 to January 1949	22	30	
February 1957 to March 1961	50	15	
March 1964 to December 1965	21	15	
Total number of months of good foreca Mean $(80 \div 4) = 20$	asts	123	80
Period of disturbed forecasts			
December 1944 to March 1947	28	45	
February 1949 to January 1957	96	42	
March 1961 to February 1964	36	40	
Total number of months of		_	
disturbed forecasts		160	127
Total months of forecasts Mean $(127 \div 3) = 42$		283	

Volcanoes and bombs produce similar effects. In Volume 4 of the *Annals of the Smithsonian Astrophysical Observatory*, page 128, the effect of the eruption of the volcano Katmai, 6 June 1912, is recorded in tabulating the multitude of pyrheliometer and bolometric-spectral observations made at Mount Wilson in 1912, 1913, and 1914. With a general mean of 1.370 calories in pyrheliometer readings, the departures noted in Augusts of 1912, 1913, and 1914 were as follows:

1912.		4	۰	۰	٥	9	۵	u		239
1913.				a		a	0	0	a	101
1914.										022

So it is not remarkable to find that, in various intervals of months and years, large discrepancies between long-range forecasts (based on normal atmospheric conditions) occurred after the bombing between 1944 and 1960.

APPENDIX

Publ. No.	Date	Smithsonian Miscellaneous Collections	Title and Author
2825	1925	Volume 77, No. 5	Solar variation and forecasting, C. G. Abbot
3114	1931	Volume 85, No. 1	Weather dominated by solar changes, C. G. Abbot
3339	1935	Volume 94, No. 10	Solar radiation and weather studies, C. G. Abbot
3637	1941	Volume 101, No. 1	An important weather element hitherto generally disregarded, C. G. Abbot
3641	1941	Volume 101, No. 5	On solar-constant and atmospheric temperature changes, Henryk Arctowski
3765	1944	Volume 104, No. 3	A 27-day period in Washington pre- cipitation, C. G. Abbot
3771	1944	Volume 104, No. 5	Weather predetermined by solar variation, C. G. Abbot
3807	1945	Volume 104, No. 13	Correlations of solar variation with Washington weather, C. G. Abbot
3893	1947	Volume 107, No. 4	The sun's short regular variation and its large effect on terrestrial temperatures, C. G. Abbot
3901	1947	Volume 107, No. 9	Precipitation affected by solar varia- tion, C. G. Abbot
3916	1948	Volume 110, No. 1	Solar variation attending West Indian hurricanes, C. G. Abbot
3940	1948	Volume 110, No. 6	Magnetic storms, solar radiation, and Washington temperature changes, C. G. Abbot
3990	1949	Volume 111, No. 13	Short periodic variations and the tem- peratures of Washington and New York, C. G. Abbot
4015	1950	Volume 111, No. 17	Periodic influence of Washington and New York weather of 1949 and 1950, C. G. Abbot
4088	1952	Volume 117, No. 10	Periodicities in the solar-constant measures, C. G. Abbot
4090	1952	Volume 117, No. 11	Important interferences with normals in weather records, associated with sunspot frequency, C. G. Abbot
4103	1953	Volume 121, No. 5	Solar variation and precipitation at Albany, N.Y., C. G. Abbot
4135	1953	Volume 122, No. 4	Solar variation a leading weather ele- ment. C. G. Abbot
4211	1955	Volume 128, No. 3	Sixty-year weather forecasts, C. G. Abbot
4213	1955	Volume 128, No. 4	Periodic solar variation, C. G. Abbot

Publ.	Date	Smithsonian Miscellaneous Collections	Title and Author
4222	1955	Volume 131, No. 1	Leading operations of the Smithsonian Astrophysical Observatory, 1895- 1955, C. G. Abbot
4265	1956	Volume 134, No. 1	Periods related to 273 months or 223/4 years, C. G. Abbot
4338	1958	Volume 135, No. 10	Periodicities in ionospheric data, C. G. Abbot
4352	1959	Volume 138, No. 3	Long-range weather forecasting. C. G. Abbot
4390	1960	Volume 139, No. 9	A long-range forecast of United States precipitation. C. G. Abbot
4462	1961	Volume 143, No. 2	Sixteen-day weather forecasts from satellite observations, C. G. Abbot
4471	1961	Volume 143, No. 5	A long-range temperature forecast. C. G. Abbot
4545	1963	Volume 146, No. 3	Solar variations and weather, C. G. Abbot
4656	1966	Volume 148, No. 7	An account of the Smithsonian Insti- tution, 1904-1953, C. G. Abbot
4659	1966	Volume 148, No. 8	Forecasting from harmonic periods in precipitation, C. G. Abbot
4694	1967	Volume 151, No. 5	Precipitation in five countries, C. G. Abbot
4711	1967	Volume 152, No. 5	Supplement to a long-range forecast of United States precipitation, C. G.
			Abbot
4722	1967	Volume 152, No. 6	Solar magnetism and world weather. C. G. Abbot

CHARLES AND AND AND ADDRESS OF THE A





